

Title page

Title: Factors associated with adherence to a Mediterranean diet in adolescents from the northern region of Spain.

Author names and affiliations:

Raúl Jiménez Boraita ¹
University of La Rioja, Logroño, Spain
Master's Degree
rauljbcity@gmail.com

Daniel Arriscado Alsina ¹
University of La Rioja, Logroño, Spain
Doctoral Degree
danielarriscado@hotmail.com

Esther Gargallo Ibor ¹
University of La Rioja, Logroño, Spain
Doctoral Degree
esther.gargallo@unirioja.es

Josep María Dalmau Torres ¹
University of La Rioja, Logroño, Spain
Doctoral Degree
jose-maria.dalmau@unirioja.es

¹ Department of Educational Sciences, University of La Rioja, C/ Luis de Ulloa s/n, Edificio Vives, CP 26002 Logroño, Spain

Corresponding author:

Raúl Jiménez Boraita, Avenida Fuenmayor nº5 1ºD 26350 Cenicero, La Rioja Spain,
tf. +34 620484910, e-mail: rauljbcity@gmail.com

Acknowledgements

The authors would like to thank all students, parents and teachers of the schools that participated in this study.



This peer-reviewed article has been accepted for publication but not yet copyedited or typeset, and so may be subject to change during the production process. The article is considered published and may be cited using its DOI

10.1017/S0007114520003803

The British Journal of Nutrition is published by Cambridge University Press on behalf of The Nutrition Society

Funding

The study was partially financed by the Institute of Riojan Studies (IER) of the Government of La Rioja through “Resolución nº 55/2018, de 9 de julio, de la gerencia del instituto de estudios riojanos para la concesión de ayudas para estudios científicos de temática riojana convocadas para el año 2018-2019.”

Author Contributions

All authors were involved in study design, data analysis, data interpretation, literature search and writing the paper and had final approval of the submitted and published versions. None of the authors have any competing interests in the manuscript

Conflict of interest

The authors declare no potential conflicts of interest.

Ethical standard

Written informed consent was requested from participants’ parents or legal guardians. Collaboration of adolescents in the present study was voluntary and all participants provided verbal consent. Fundamental ethics of the Declaration of Helsinki were respected. In addition, the project was approved by the Ethical Committee for Clinical Research of La Rioja.

Title: Factors associated with adherence to a Mediterranean Diet in adolescents from La Rioja (Spain).

ABSTRACT

The aim of the present study was to describe Mediterranean diet adherence within a population of adolescents and to analyse the association of multiple factors with adherence. This included a consideration of diverse physical and mental health indicators. The present study was conducted with a representative sample of 761 adolescents (14.51 ± 1.63 years) from 25 educational centres in a northern region of Spain. Mediterranean diet adherence was evaluated, alongside their health-related quality of life, self-esteem, body image satisfaction, body mass index, physical activity level, maximum oxygen consumption, hours of nightly sleep, sociodemographic factors and academic performance. Forty-nine percent of the adolescent population reported high Mediterranean diet adherence. The female gender and higher levels of physical activity were found to be predictive factors of adherence to the Mediterranean diet. In addition, maximum oxygen consumption, the presence of environments favourable towards physical activity engagement and higher self-esteem were also predictive in females, whilst better academic performance and more nightly sleep were additional predictors in males. The associations found between Mediterranean diet, and other health indicators and habits, highlight the need to develop promotion strategies from an inter-disciplinary and transversal standpoint.

Keywords: Mediterranean diet, adolescent, habits, health.

INTRODUCTION

The MD is considered to be one of the most influential dietary patterns in relation to health. Its composition is principally based on food groups that are characteristic of Mediterranean zones. At the same time, it also incorporates a way of cooking and a lifestyle that includes physical activity engagement, sufficient rest and socialisation during meals. The MD possesses common patterns such as the inclusion of cereals, vegetables and fruits in all meals, together with a moderate consumption of dairy products. Equally, the use of olive oil as a main source of fat is considered fundamental and weekly incorporation of animal or vegetable protein is also required. On the one hand, with regards to proteins that are of animal origin, consumption preferences should be directed towards fish (≥ 2), eggs (2-4 times a week) and white meats (≥ 2), with red meats being consumed in lower quantities and at a lower frequency (≤ 2). On the other hand, legumes are a main source of vegetable protein (≥ 2). Finally, water provides the main source of hydration⁽¹⁾.

MD adherence has been demonstrated to exert protective effects against cardiovascular diseases, coronary diseases, acute myocardial infarctions and cancer^(2, 3). Relations have been found in adolescents between a healthy diet and various cardiovascular risk factors, such as blood pressure⁽⁴⁾ and cholesterol⁽⁵⁾. On the other hand, the MD has also been demonstrated to exert positive effects on quality of life perceptions, physical activity levels (PA) and sleep quality. Further, impact has been shown on obesity, although these results are not conclusive^(6, 7, 8). In addition, greater MD adherence is associated with a lower risk of suffering from mental illnesses, especially those with a depressive symptomology⁽⁹⁾.

Nonetheless, despite the diverse array of benefits shown for health, trends in recent years show a reduction in adherence to the traditional MD, and a rise in less healthy habits amongst children and adolescents^(10, 11). In this sense, maintenance of healthy dietary patterns during adolescence could prevent the risk of suffering from health problems during adulthood^(12, 13, 14).

Thus, evaluation of the factors influencing MD adherence is key for better understanding the current situation. In this way, choices regarding food consumption can be determined at an individual level according to age, nationality, sex, preferences, nutritional knowledge, physiological state and psychological factors^(15, 16, 17). On the other hand, sociocultural factors such as social norms, the influence of family and peers, socioeconomic status, the media and social networks appear to play a mediatory role over dietary habits^(18, 19). Finally, the influence of setting, in the same way as access to and availability of healthy foods, also intervenes in dietary choices⁽²⁰⁾.

The majority of studies dealing with MD in adolescents studied its relationship with obesity or other variables mainly associated with physical health. However, the present study approaches the analysis of adherence to a MD from a broader perspective, examining its relationship with various indicators, not only of physical health but also psychological and social health, whilst also analysing its association with different sociodemographic variables and potential predictive factors. To this end, various instruments were used to assess the different dimensions of health. Instruments assessed adolescents' adherence to the Mediterranean diet, physical activity level,

health-related quality of life, self-esteem, body image satisfaction, hours of nightly sleep, body mass index, maximal oxygen uptake, academic performance and sociodemographic factors.

METHODS AND PROCEDURES

Participants

A descriptive, cross-sectional and correlational study was carried out with a representative sample of adolescents undertaking secondary education at educational centres in La Rioja, a region in the north of Spain. The sample was obtained through stratified random sampling according to clusters, in which secondary education classrooms in all centres of the region provided the sampling unit. Of a total of 6.018 adolescents enrolled in these classrooms, it was estimated that the number required for sampling to be representative was 362 (95% confidence interval). A final sample of 761 adolescents was obtained from 25 educational centres. Participants were aged between 12 and 17 years (14.51 ± 1.63 years), with 49.7% being female and 51.3% being male.

Procedure

Written informed consent was requested from adolescents' parents or legal guardians. Collaboration of adolescents in the present study was voluntary and all provided verbal consent. Fundamental ethics of the Declaration of Helsinki were respected. In addition, the project was approved by the Ethical Committee for Clinical Research of La Rioja. Members of the research team carried out field work during teaching hours at the educational centres, with all members following the same action protocol: Self-completion questionnaire, anthropometric measures and physical aptitude test.

Instruments

Mediterranean diet (MD) adherence was evaluated using the KIDMED questionnaire developed by Serra et al.⁽²¹⁾. The instrument comprises sixteen items with dichotomous response options ("yes" or "no") in relation to the consumption of foods related with Mediterranean dietary patterns. The procedure established by the author was followed in order to derive an overall adherence score, with possible values falling between minus four and twelve. The final derived score was then classified according to the parameters indicated by the author of the questionnaire: low (≤ 3), medium (4-7) or high (≥ 8). For data handling, adolescents reporting low (4.7%) and medium (46.3%) MD adherence were grouped together due to the low percentage reporting the former.

Health-related quality of life (HRQL) was estimated through the KIDSCREEN-27 questionnaire, developed and validated within Spanish adolescents by Aymerich et al.⁽²²⁾. This questionnaire is formed by 27 items which are grouped according to five dimensions and rated along a Likert scale. The dimensions are as follows: Physical wellbeing, psychological wellbeing, autonomy and parenting, school environment, and social support and friends. Obtained data were handled in accordance with the parameters established by the authors of the questionnaire, with higher values corresponding to more positive perceptions of HRQL⁽²³⁾.

Evaluation of *self-esteem* was performed using Rosenberg's self-esteem scale, adapted and validated by Atienza et al.⁽²⁴⁾ in Spanish adolescents. This scale is formed by ten items, with responses being rated from one to four, permitting final scores of between 10 and 40. In this way, higher scores are related with higher self-esteem.

For evaluation of *perceptions of and satisfaction with body image*, the method described by Stunkard and Stellar⁽²⁵⁾ and adapted by Marrodán et al.⁽²⁶⁾ was used. For this evaluation, nine silhouettes of female figures are used in addition to nine male figures. The silhouettes show a progression towards a stockier appearance and correspond to different BMI values, running from 17kg/m² up to 33kg/m². Participants were asked to choose two silhouettes, the one with which they most identify themselves based on their actual appearance, and the silhouette representing the figure they wished they had. In this way, the difference between both silhouettes is calculated, grouping adolescents according to two categories: Adolescents who are satisfied with their body image (values between -2 and 2) and those who are dissatisfied (all values outside of this range).

Physical activity (PA) level was estimated using the Physical Activity Questionnaire for Adolescents (PAQ-A), adapted and validated within Spanish adolescents by Martínez et al.⁽²⁷⁾. This questionnaire evaluates the PA engaged in during the last seven days, as a function of frequency and time-frames. Obtained values are between one and five, with higher values being associated with greater engagement in PA. In addition, two questions are posed in relation to engagement in organised sporting activities and the mode of transport used to arrive at the educational centre: "Do you engage in any extra-curricular sporting activity after school?" and "Do you exercise when you go from home to school (walking, cycling, skating...)" With the aim of calculating hours of nightly sleep, adolescents were also asked about the time at which they typically went to bed and woke up.

Evaluation of *setting* was analysed through the ALPHA environmental questionnaire developed by Spittaels et al.⁽²⁸⁾ within the ALPHA Project of the European framework. This questionnaire evaluates perceptions of factors within one's immediate environment (from within approximately 1.5 km of the home) which may influence PA engagement. A version of this instrument was used which was adapted for the Spanish youth population and validated by García-Cervantes et al.⁽²⁹⁾. Scores are obtained by summing together the ten questionnaire items. Higher results represent a more favourable setting for engaging in PA. Results were categorised using the median as the cut-point. This enabled two potential settings: Favourable or unfavourable.

With regards to sociodemographic data, adolescents reported their sex, date of birth and nationality ("born in Spain" or "born in another country"). For evaluation of *socioeconomic status*, the family affluence scale (FAS II) was used⁽³⁰⁾. This constitutes four questions related with family possession of material goods. The final score ranges between zero and nine, with results being categorised in the following way: Low (≤ 2), medium (3-5) or high (≥ 6). For data handling, adolescents reporting a low socioeconomic status (1.8%) and a medium socioeconomic status (28%) were grouped together due to the low percentage reporting the former.

Likewise, the *Oviedo response frequency scale* (INF-OV)⁽³¹⁾ was also used with the aim of detecting and excluding from data analysis any questionnaire data which had been provided in a random, dishonest or pseudo-random way. The INF-OV is a self-report instrument which includes questions which demand dichotomous (yes or

no) responses. In this way, six items were introduced in an interspersed way throughout the questionnaire (for example, “have you ever used the bus?” or “have you ever seen children playing in the park?”). Adolescents who provided responses that most contradicted logic were excluded from the analysis, more specifically, two were excluded for this reason.

For the measurement of height and weight a Holtain® (Holtain Ltd., Dyfed, United Kingdom) stadiometer and SECA® scale (713, Hamburg, Germany) were used, with a precision of one millimetre and 0.1 kg, respectively. Anthropometric measurements were performed according to the protocols of the International Society for the Advancement of Kinanthropometry (ISAK)⁽³²⁾. Following this, BMI was calculated and adolescents were categorised as a function of reference values established by the WHO: Normal weight, overweight and obesity⁽³³⁾.

Evaluation of *aerobic capacity* was conducted with the Course-Navette Test. For this, two transversal lines marked a distance of 20 metres, indicating the start and finish of the run route. Participating adolescents had to maintain a race speed that was dictated by an acoustic signal. This determined the time in which participants had to successively cover the distance between both lines. The acoustic signal began the race at a speed of 8.5 km/h, increasing by 0.5 km/h every minute. In this way, each adolescent finished the test when they stopped or failed, on two successive occasions, to complete the route within the time indicated by the signal. Through the results obtained, *maximal oxygen consumption* ($VO_2^{m\acute{a}x}$) was calculated according to the formula stipulated by the author who conceived the test⁽³⁴⁾.

Finally, *academic performance* was analysed through the grades provided by the Educational Council of the Government of La Rioja. Specifically, this outcome was obtained by recording the average mark obtained by the adolescent for the school year they were undertaking at the time of the study. Ninety percent of families gave their express consent for these grades to be provided to the research team.

Statistical analysis

Quantitative variables were represented according to means and standard deviations, whilst in exchange, qualitative variables are given according to frequencies. Normality and homoscedasticity of the data were verified with the Kolmogorov-Smirnov test and the Levene test, respectively. Means were compared through the Student’s t-test and the Mann-Whitney U for normally and non-normally distributed variables, respectively. The Pearson chi-squared test was used to analyse the associations between qualitative variables.

Factors associated with the MD score were analysed via univariate analysis. Pearson and Spearman indices were used for normally and non-normally distributed variables, respectively. Following this, a multiple linear regression model was developed using backward elimination. The aim of this was to identify predictors of the MD score and to control for the effect of confounding variables. This analysis included the variables of age, nationality, sex, socioeconomic status, setting, body mass index, body satisfaction, health-related quality of life, self-esteem, physical activity, maximal oxygen consumption, hours of nightly sleep and academic performance. This same analysis was carried out as a function of sex. Models met the assumptions of multiple regression in terms of linearity, homoscedasticity, normality, independence and non-multicollinearity. All statistical analyses were carried

out using the program IBM-SPSS® in its version 25 for Windows. Statistical significance was established at $p < 0.05$.

RESULTS

Adolescents' age, health-related quality of life, self-esteem, physical activity engagement, aerobic capacity, hours of nightly sleep and body mass index are shown in Table 1 as a function of MD adherence. For all variables, analysis revealed significantly higher values on the part of adolescents with high MD adherence. The only exceptions were found for age, where a significant inverse relationship was uncovered, and BMI, where no significant differences were found.

Table 2 presents outcomes for the analysis of influential factors with regards to MD adherence. Greater adherence is observed on the part of adolescents with a high socioeconomic status and who reside within settings that are favourable towards PA engagement. In the same way, greater adherence was reported by adolescents who were satisfied with their body image. Despite failing to find statistically significant findings in relation to nationality, a greater prevalence of high adherence was highlighted amongst natives (50.1%), relative to migrants (40.1%).

Table 3 shows the frequency with which certain foods are consumed as a function of various sociodemographic factors. Migrants reported a lower consumption than natives of dairy, fish and legumes, in addition to a higher consumption of pasta or rice and sweets. With respect to socioeconomic status, adolescents with a high status reported significantly higher percentages for the consumption of breakfast cereals, fruit and juices, fish, legumes and second servings of fruit, and dairy. In contrast, consumption of pasta, rice and sweets was higher in adolescents with a low/medium socioeconomic status. Finally, differences were also observed as a function of setting in that, those who were residents in zones that favoured engagement in PA, reported higher frequencies for the consumption of fruit and juices, and vegetables and fish, alongside a lower consumption of pasta and rice.

Table 4 details the correlation coefficients relating MD with age, socioeconomic status, sleep, HRQL, self-esteem, PA, $VO_2^{m\acute{a}x}$ and academic performance. Higher MD adherence was inversely associated with age and positively associated with socioeconomic status, hours of nightly sleep, HRQL, self-esteem, PA engagement, aerobic capacity and academic performance. This association was present in both females and males.

Finally, the multiple regression model using backward elimination is shown in Table 5. The female gender, having a higher level of PA and self-esteem, better academic performance, and achieving more hours of nightly sleep were found to be predictive factors of MD adherence. When considering gender-related differences, better academic performance and a greater number of hours of nightly sleep were only predictors in the case of boys, whilst $VO_2^{m\acute{a}x}$, self-esteem and environment were only predictors for girls.

DISCUSSION

Results of the present study showed that adherence to a MD was related to various physical and mental aspects including sociodemographic variables, lifestyle habits and health indicators. Through scores obtained on the

KIDMED, we can see that 49% of adolescents presented high values of MD adherence, 46.3% had moderate adherence and 4.7% showed a low adherence. In this sense, results show similar trends to those reported in the Enkid study by Serra-Majem et al.⁽²¹⁾ with a representative sample of the Spanish population aged from 2 to 24 years (with 46.4% reporting high adherence). It also corroborates results obtained in a study by Mariscal-Arcas et al.⁽³⁵⁾ in adolescents from Granada (46.9%). Nonetheless, values for high adherence were higher than those found in a sample of Andalusians (30.9%) and Catalonians (26%)^(36, 37), and were relatively higher than those found in other homologous Mediterranean countries such as Italy, Greece and Turkey (9.1%, 26% and 22.9%, respectively)^(38, 39, 40). Such differences, identified as a function of geographic region, underline the importance of better understanding the factors that impact upon MD adherence.

Through multivariate analysis, we were able to establish predictors of MD, with gender emerging as a determining factor. Along the same lines, a study carried out in 21 European countries revealed that females were more likely to consume fruits and vegetables⁽⁴¹⁾. This finding could be partly explained by the greater interest shown by females in weight control and body image⁽⁴²⁾. In this way, pressure resulting from the promotion of beauty ideals of thinness by the mass media and social media encourages healthy food choices, leading to the avoidance of fat and salt and a higher consumption of fruit and fiber⁽⁴³⁾.

With regards to PA, engagement was significantly higher in adolescents with MD high adherence, with this proving to be a predictive factor in both sexes. Thus, a higher level of PA has been associated with a greater consumption of fish, fruit, vegetables, nuts, legumes and olive oil⁽⁴⁴⁾. Further, in agreement with the present study, the aforementioned research also demonstrated that adolescents with high MD adherence reported significantly higher values in relation to VO₂máx. This finding may be due to PA levels themselves, with numerous studies unveiling the association between PA and subsequent physical condition⁽⁴⁵⁾. However, our results only showed the predictive value of VO₂max in the case of girls.

With regards to setting, adolescents residing in favourable contexts for PA engagement had significantly higher adherence rates to the MD, alongside a higher consumption of fruit, juices, vegetables and fish. This fact could be associated with socioeconomic status since present results confirmed a positive correlation between setting and socioeconomic status ($r = 135$). In this way, adolescents living in areas with greater opportunities for PA also had a higher socioeconomic status, with this potentially having a direct impact on access to food purchases. However, our results identified the setting as a predictor only in the case of girls.

In the same way, self-esteem was also shown to be a valuable predictor. Previous studies have shown that the consumption of fruits and vegetables is directly associated with psychological health⁽⁴⁶⁾, exercising a protective role against depression, anxiety and psychological distress⁽⁴⁷⁾. However, our results only showed the predictive value of self-esteem on MD in girls. In this sense, the importance of body image on girls' self-esteem could lead to a desire for weight loss and, therefore, greater diet control. In this way, improvements to self-esteem appear to be key for preventing the development of eating disorders in girls⁽⁴⁸⁾.

In respect to hours of nightly sleep, significant differences were also found, with adolescents who more strictly adhered to a MD reporting more hours of sleep. These results collaborate those reported by Rosi et al.⁽⁷⁾. In the same way, Ferranti et al.⁽⁸⁾ reported that sleep duration was positively associated with the intake of fruits and vegetables, and inversely related with the consumption of sweets and snacks. In our results, this duration turned out to be a predictor of MD in the case of boys.

Finally, academic performance was superior within adolescents with better MD adherence. This relationship has already been reported previously and has been determined to occur independently of factors such as BMI and PA⁽⁴⁸⁾. In the same way, Frisardi et al.⁽⁴⁹⁾ confirmed the relationship between consumption of foods inherent to the MD and conservation and improvement of cognitive processes. Such dietary practices appear to offer protection through vascular inflammation and oxidation which results in better neuronal conservation. Along the same lines, Chacón-Cuberos et al.⁽⁵⁰⁾ uncovered positive associations between high MD adherence and processes related with learning strategies and motivation. This led to better scores for elaboration strategies, organisation, critical thinking, effort regulation, and study habits and time. In our study, academic performance was even a predictor of MD adherence in boys. In this way, these results could serve as a call to promote the importance of a healthy diet.

Besides the predictive factors described above, MD was shown to be associated with other sociodemographic variables and indicators related with adolescent's mental wellbeing. Firstly, individuals with high NSE reported greater adherence to Mediterranean dietary patterns. This coincides with studies that have evidenced greater consumption of fruit, fish and legumes, in addition to lower consumption of pasta, rice and sweets⁽²¹⁾. On the other hand, MD adherence was inversely correlated with age. This could be associated with the greater influence of peers with increasing age, alongside increases in autonomy and financial independence⁽¹⁸⁾. With regards to indicators of mental wellbeing, adolescents with greater MD adherence also reported higher levels of CVRS and body satisfaction. In this sense, previously conducted studies have already uncovered relationships between the MD and various quality of life dimensions⁽⁵¹⁾, whilst also revealing dietary restrictions which are non-characteristics of Mediterranean patterns amongst adolescents who are unhappy with their body weight⁽⁵²⁾. Nonetheless, none of these associations proved to be determinant in the multivariate analysis. Thus, given the potentially confusing effects produced, present results should be treated cautiously.

One of the main strengths of the present study is based on its inclusion of a representative sample which enables us to extrapolate the influence of sociodemographic factors on MD adherence to wider populations. This is also true for the associations between these factors and diverse health indicators, both physical and mental, providing a global view in the analysis of MD. Nonetheless, the present study is not exempt from limitations. Self-

complete questionnaires were used which are susceptible to the subjectivity of participants. Future studies should be considered which use instruments that provide greater objectivity and data precision, including food consumption diaries or accelerometers. This being said, the data collection protocol was the same for all adolescents, protected anonymity and used valid instruments which had demonstrable reliability within populations similar to that used in the present study. On the other hand, use of a cross-sectional design in the present study makes it impossible to determine the direction of associations. This makes it necessary to develop future research studies which are longitudinal and experimental in nature.

CONCLUSION

The results obtained show that 49% of the included adolescent population presented high MD adherence. The female gender, higher PA levels and self-esteem, better academic performance, and more hours of nightly sleep were found to be predictive factors of MD adherence. When considering gender-related differences, better academic performance and a greater number of hours of nightly sleep were predictors only in the case of boys, whilst VO₂max, self-esteem and environment were predictors for girls. The relationships found between MD adherence and other indicators and habits highlight the need to develop promotional strategies from an interdisciplinary and transversal standpoint. Given the tendency for adolescents move away from the MD as they age, educational centres and related institutions are advised to promote healthy dietary patterns based on the MD.

REFERENCES

1. Bach-Faig A, Berry EM, Lairon D, *et al.* (2011) Mediterranean diet pyramid today. Science and cultural updates. *Public health nutrition* 14, 2274-2284.
2. Rosato V, Temple NJ, La Vecchia C, *et al.* (2019) Mediterranean diet and cardiovascular disease: A systematic review and meta-analysis of observational studies. *European Journal of Nutrition* 58, 173–191.
3. Schwingshackl L, Schwedhelm C, Galbete C, *et al.* (2017) Adherence to Mediterranean diet and risk of cancer: an updated systematic review and meta-analysis. *Nutrients* 9,1063.
4. Bacopoulou F, Landis G, Rentoumis A, *et al.* (2017) Mediterranean diet decreases adolescent waist circumference. *European journal of clinical investigation* 47, 447-455.
5. Zhong VW, Lamichhane AP, Crandell JL *et al.* (2016) Association of adherence to a Mediterranean diet with glycemic control and cardiovascular risk factors in youth with type I diabetes: the SEARCH Nutrition Ancillary Study. *European journal of clinical nutrition* 70, 802.
6. Evaristo OS, Moreira C, Lopes L, *et al.* (2018) Associations between physical fitness and adherence to the Mediterranean diet with health-related quality of life in adolescents: Results from the LabMed Physical Activity Study. *The European Journal of Public Health* 28, 631-635.
7. Rosi A, Giopp F, Milioli G, *et al.* (2020) Weight Status, Adherence to the Mediterranean Diet, Physical Activity Level, and Sleep Behavior of Italian Junior High School Adolescents. *Nutrients* 12, 478.
8. Ferranti R, Marventano S, Castellano S, *et al.* (2016) Sleep quality and duration is related with diet and obesity in young adolescent living in Sicily, Southern Italy. *Sleep Science* 9, 117-122.
9. Altun A, Brown H, Szoeki C, *et al.* (2019) The Mediterranean dietary pattern and depression risk: A systematic review. *Neurology, Psychiatry and Brain Research* 33, 1-10.
10. Cabrera SG, Fernández NH, Hernández CR, *et al.* (2015) KIDMED test; prevalence of low adherence to the Mediterranean Diet in children and young; a systematic review. *Nutrición hospitalaria* 32, 2390-2399.
11. Thana YA, Takruri HR, Tayyem RF (2019) Dietary Practices and Nutrient Intake among Adolescents: A General Review. *Obesity Medicine*, 100145.
12. Hu T, Jacobs Jr DR, Larson NI, *et al.* (2016) Higher diet quality in adolescence and dietary improvements are related to less weight gain during the transition from adolescence to adulthood. *The Journal of pediatrics* 178, 188-193.

13. Wennberg M, Gustafsson PE, Wennberg P, *et al.* (2016) Irregular eating of meals in adolescence and the metabolic syndrome in adulthood: results from a 27-year prospective cohort. *Public health nutrition* 19, 667-673.
14. Pearson CM, Miller J, Ackard DM, *et al.* (2017) Stability and change in patterns of eating disorder symptoms from adolescence to young adulthood. *International Journal of Eating Disorders* 50, 748-757.
15. Arcila-Agudelo AM, Ferrer-Svoboda C, Torres-Fernández T, *et al.* (2019) Determinants of Adherence to Healthy Eating Patterns in a Population of Children and Adolescents: Evidence on the Mediterranean Diet in the City of Mataró (Catalonia, Spain). *Nutrients* 11, 854.
16. Tarabashkina L, Quester P, Crouch R (2016) Exploring the moderating effect of children's nutritional knowledge on the relationship between product evaluations and food choice. *Social Science & Medicine* 149, 145-152.
17. Conner M, Povey R, Sparks P, *et al.* (1998) Understanding dietary choice and dietary change: contributions from social psychology. In *The Nation's Diet: The Social Science of Food Choice* [Murcott A, editor]. London: Longman:
18. Moreno-Maldonado C, Ramos P, Moreno C, *et al.* (2018) How family socioeconomic status, peer behaviors, and school-based intervention on healthy habits influence adolescent eating behaviors. *School Psychology International* 39, 92-118.
19. Higgs S & Thomas J (2016) Social influences on eating. *Current Opinion in Behavioral Sciences* 9, 1-6.
20. Gustafson A, Jilcott S, McDonald J, *et al.* (2017) Direct effects of the home, school, and consumer food environments on the association between food purchasing patterns and dietary intake among rural adolescents in Kentucky and North Carolina, 2017. *International journal of environmental research and public health* 14, 1255.
21. Serra L, Ribas L, Ngo J, *et al.* (2004) Food, youth and the Mediterranean diet in Spain. Development of KIDMED, Mediterranean Diet Quality Index in children and adolescents. *Public health nutrition* 7, 931-935.
22. Aymerich M, Berra S, Guillamón I, *et al.* (2005) Desarrollo de la versión en español del KIDSCREEN: un cuestionario de calidad de vida para la población infantil y adolescente. *Gaceta Sanitaria* 19, 93-102
23. The KIDSCREEN Group Europe (2006) *The KIDSCREEN questionnaires- quality of life questionnaires for children and adolescents*. Lengerich: Pabst Science Publishers.

24. Atienza FL, Moreno Y, Balaguer I (2000) Análisis de la dimensionalidad de la Escala de Autoestima de Rosenberg en una muestra de adolescentes valencianos. *Revista de Psicología Universitat Tarraconensis* 22,29-42.
25. Stunkard A & Stellar E (1990) Eating and its disorders. In *Body Images*, pp. 3-20 [Cash T and Pruzinsky T, editors]. Nueva York: Guilford Press,
26. Marrodan V, Montero MS, Mesa JL, *et al.* (2008) Realidad, percepción y atractivo de la imagen corporal: condicionantes biológicos y socioculturales. *Zainak, Cuadernos de Antropología-Etnografía* 30, 15-28.
27. Martínez D, Martínez de Haro V, Pozo T, *et al.* (2009) Fiabilidad y validez del cuestionario de actividad física PAQ-A en adolescentes españoles. *Revista española de salud pública* 83, 427-439.
28. Spittaels H, Verloigne M, Gidlow C, *et al.* (2010) Measuring physical activity-related environmental factors: reliability and predictive validity of the European environmental questionnaire ALPHA. *International Journal of Behavioral Nutrition and Physical Activity* 7, 48.
29. García-Cervantes L, Martínez-Gómez D, Rodríguez-Romo G, *et al.* (2014) Reliability and validity of an adapted version of the ALPHA environmental questionnaire on physical activity in Spanish youth. *Nutrición hospitalaria* 30, 1118-1124.
30. Currie C, Molcho M, Boyce W, *et al.* (2008) Researching health inequalities in adolescents: the development of the Health Behaviour in School-Aged Children (HBSC) family affluence scale. *Social science & medicine* 66, 1429-1436.
31. Fonseca E, Paño M, Lemos S, *et al.* (2009) Validation of the Schizotypal Personality Questionnaire Brief form in adolescents. *Schizophrenia Research* 111, 53-60.
32. Stewart A, Marfell-Jones M, Olds T, *et al.* (2011) *International standards for anthropometric assessment*. New Zealand: Lower Hutt;2011.
33. Onis MD, Onyango AW, Borghi E, *et al.* (2007) Development of a WHO growth reference for school-aged children and adolescents. *Bulletin of the World health Organization* 85, 660-667.
34. Leger LA, Mercier D, Gadoury C, *et al.* (1988) The multistage 20 metre shuttle run test for aerobic fitness. *Journal of sports sciences* 6, 93-101.
35. Mariscal-Arcas M, Rivas A, Velasco J, *et al.* (2009) Evaluation of the Mediterranean Diet Quality Index (KIDMED) in children and adolescents in Southern Spain. *Public health nutrition* 12, 1408-1412.
36. Grao-Cruces A, Nuviala A, Fernández-Martínez A, *et al.* (2013) Adherence to the Mediterranean diet in rural and urban adolescents of southern Spain, life satisfaction, anthropometry, and physical and sedentary activities. *Nutrición hospitalaria* 28, 1129-1135.

37. Fauquet J, Sofi F, López-Guimerà G, *et al.* (2016) Mediterranean diet adherence among Catalanian adolescents: socio-economic and lifestyle factors. *Nutrición hospitalaria* 33, 1283-1290.
38. Mistretta A, Marventano S, Antoci M, *et al.* (2017) Mediterranean diet adherence and body composition among Southern Italian adolescents. *Obesity research & clinical practice* 11, 215-226.
39. Vassiloudis I, Yiannakouris N, Panagiotakos DB, *et al.* (2017) Adherence to the Mediterranean diet and specific lifestyle habits are associated with academic performance in Greek adolescents. *Mediterranean Journal of Nutrition and Metabolism* 10, 93-103.
40. Sahingoz SA & Sanlier N (2011) Compliance with Mediterranean Diet Quality Index (KIDMED) and nutrition knowledge levels in adolescents. A case study from Turkey. *Appetite* 57, 272-277.
41. Stea TH, Nordheim O, Bere E, *et al.* (2020) Fruit and vegetable consumption in Europe according to gender, educational attainment and regional affiliation—A cross-sectional study in 21 European countries. *Plos one* 15(5), e0232521.
42. Field AE, Camargo CA, Taylor CB, *et al.* (2001) Peer, parent, and media influences on the development of weight concerns and frequent dieting among preadolescent and adolescent girls and boys. *Pediatrics* 107(1), 54-60.
43. Wardle J, Haase AM, Steptoe A, *et al.* (2004) Gender differences in food choice: the contribution of health beliefs and dieting. *Annals of behavioral medicine* 27(2), 107-116.
44. Muros JJ, Cofre-Bolados C, Arriscado D, *et al.* (2017) Mediterranean diet adherence is associated with lifestyle, physical fitness, and mental wellness among 10-y-olds in Chile. *Nutrition* 35, 87-92.
45. DeFina LF, Haskell WL, Willis BL, *et al.* (2015) Physical activity versus cardiorespiratory fitness: two (partly) distinct components of cardiovascular health?. *Progress in cardiovascular diseases* 57, 324-329.
46. Boehm JK, Soo J, Zevon ES, *et al.* (2018) Longitudinal associations between psychological well-being and the consumption of fruits and vegetables. *Health Psychology* 37, 959.
47. Sadeghi O, Keshteli AH, Afshar H, *et al.* (2019) Adherence to Mediterranean dietary pattern is inversely associated with depression, anxiety and psychological distress. *Nutritional neuroscience* 1-12.
48. Argyrides M & Kkeli N (2015) Predictive factors of disordered eating and body image satisfaction in Cyprus. *International Journal of Eating Disorders* 48, 431-435.
49. Frisardi V, Panza F, Seripa D, *et al.* (2010) Nutraceutical properties of Mediterranean diet and cognitive decline: possible underlying mechanisms. *Journal of Alzheimer's Disease* 22, 715-740.
50. Chacón-Cuberos R, Zurita-Ortega F, Martínez-Martínez A, *et al.* (2018) Adherence to the mediterranean diet Is related to healthy habits, learning processes, and academic achievement in adolescents: a cross-sectional study. *Nutrients* 10, 1566.
51. Knox E & Muros JJ (2017) Association of lifestyle behaviours with self-esteem through health-related quality of life in Spanish adolescents. *European journal of pediatrics* 176, 621-628.
52. Bibiloni MDM, Pich J, Pons A, *et al.* (2013) Body image and eating patterns among adolescents. *BMC public health* 13, 1104.

Table 1. Characteristics of study sample according to MD adherence

	Total		MD adherence				p-value
	N=761		Low/Mid N=388		High N=373		
	Mean	SD	Mean	SD	Mean	SD	
Age(years)	14.50	1.63	14.79	1.60	14.21	1.61	<0.001
Health-related quality of life	250.11	33.18	245.83	33.63	254.56	32.15	.003
Self esteem	32.69	4.94	32.06	5.09	33.35	4.68	<0.001
Body mass index (kg/m ²)	21.01	3.26	21.1	3.40	20.91	3.11	.637
Physical activity	2.61	0.62	2.45	0.61	2.77	0.58	<0.001
VO ² máx(ml/kg/min)	44.25	6.84	43.14	6.89	45.42	6.60	<0.001
Hours of nightly sleep	8.36	0.92	8.25	0.94	8.47	0.88	<0.001
Academic performance	6.59	1.47	6.38	±1.47	6.79	0.07	.001

Table 2. Adherence to the MD according to different factors

		MD adherence				p-value
		Low/Mid		High		
		N	%	N	%	
Nationality	Natives	338	49.9%	339	50.1%	.097
	Migrants	50	59.5%	34	40.5%	
Sex	Boys	197	51.4%	186	48.6%	.802
	Girls	191	50.5%	187	49.5%	
Socioeconomic status	Low/Mid	139	61.2%	88	38.8%	<0.001
	High	249	46.6%	285	53.4%	
Setting	Favourable	180	47%	203	53%	.027
	Unfavourable	208	55%	170	45%	
Body mass index	Normal weight	292	52%	269	48%	.325
	Overweight/Obese	96	48%	104	52%	
Body satisfaction	Satisfied	141	46.2%	164	53.8%	.032
	Dissatisfied	247	54.2%	209	45.8%	

Table 3. Consumption of foods related with MD according to nationality, socioeconomic status and setting.

	Nationality		Socioeconomic status		Setting	
	Natives n=677	Migrants n=84	Low/Mid n=227	High n=534	Unfavourable n=378	Favourable n=383
	% YES	% YES	% YES	% YES	% YES	% YES
Daily breakfast	94.4	94	92.5	95.1	94.7	94
A dairy product for breakfast (yoghurt, milk, etc.)	87.6	83.3	84.6	88.2	86.5	87.7
Cereals or grains (bread, etc.) for breakfast	80.6	82.1	75.8	83*	79.1	82.5
Commercially baked goods or pastries for breakfast	20.4	17.9	19.8	20.2	20.9	19.3
Fruit or fruit juice daily	67.2	67.9	60.4	70.2**	61.4	73.1**
Second fruit juice daily	49.2	47.6	37.4	53.9***	46.3	51.7
Second serving of dairy products daily	75.2	53.6***	65.2	76**	72	73.6
Fresh or cooked vegetables daily	72.4	76.2	69.2	74.3	70.4	75.2
Fresh or cooked vegetables >1 time/day	32.6	33.3	29.5	34.1	28	37.3**
Regular fish consumption (at least 2–3 times/week)	66.6	50**	55.5	68.7***	61.4	68.1*
Nuts regularly (≥2–3/week)	48.7	38.1	48.9	47	46.3	48.8
Fast-food restaurants >1/week	19.2	22.6	19.8	19.5	21.7	17.5
Pulses >1/week	85.7	71.4**	76.2	87.5***	85.2	83
Pasta or rice almost daily (≥5/week)	37.1	53.6**	44.9	36.3*	42.9	35*
Sweets and candies several times a day	18.5	29.8*	26.4	16.9**	22.2	17.2
Olive oil usage at home	97.9	95.2	96	98.3	97.1	98.2

p < .05*; p < .01**; p < .001***

Table 4. *Correlation coefficients related to MD adherence*

	Age	Socioeconomic status	Sleep	HRQL	Self-esteem	PA	BMI	VO2máx	Academic performance
MD	-.167**	.164**	.127**	.175**	.169**	.306**	-0.007	.166**	.154**
MD Boys	-.214**	.151**	.184**	.160**	.137**	.329**	-0.029	.194**	.165**
MD Girls	-.117*	.173**	.074	.198**	.202**	.348**	0.016	.286**	.132*

p < .05*; p < .01**

Table 5. Predictors of MD adherence

MD adherence	β	SD	p-value	95% CI	R ²
Total					
Physical activity	1.015	0.127	<0.001	0.765 - 1.264	
Academic performance	0.218	0.052	<0.001	0.116 - 0.320	
Sex (Boys)	-0.404	0.152	0.008	-0.703 - -0.106	0.143
Hours of nightly sleep	0.191	0.083	0.021	0.028 - 0.354	
Self esteem	0.028	0.016	0.073	-0.003 - 0.060	
Boys					
Physical activity	1.047	0.169	<0.001	0.715-1.379	
Academic performance	0.262	0.070	<0.001	0.125-0.400	0.170
Hours of nightly sleep	0.341	0.110	0.002	0.125-0.557	
Girls					
Physical activity	0.786	0.205	<0.001	0.382-1.189	
Maximal oxygen consumption	0.054	0.024	0.024	0.007-0.100	0.134
Self esteem	0.053	0.021	0.014	0.011-0.095	
Setting (Favourable)	0.439	0.213	0.040	0.021-0.858	

β : regression coefficient. 95% CI: 95% Confidence interval. Variables entered into model: Age, nationality, sex, socioeconomic status, setting, body mass index, body satisfaction, health-related quality of life, self-esteem, physical activity, maximal oxygen consumption, hours of nightly sleep and academic performance.